

A FORMULA FOR THE EULER CHARACTERISTICS OF EVEN DIMENSIONAL TRIANGULATED MANIFOLDS

TOSHIYUKI AKITA

A finite simplicial complex K is called an *Eulerian manifold* (or a *semi-Eulerian complex* in the literature) if all of maximal faces have the same dimension and, for every nonempty face $\sigma \in K$,

$$\chi(\text{Lk } \sigma) = \chi(S^{\dim K - \dim \sigma - 1})$$

holds, where $\text{Lk } \sigma$ is the link of σ in K and S^n is the n -dimensional sphere. Note that K is not necessary connected. Any triangulation of a closed manifold is an Eulerian manifold. More generally, a triangulation of a homology manifold without boundary provides an Eulerian manifold. The purpose of this short note is to prove the following alternative formula for the Euler characteristics of even dimensional Eulerian manifolds.

Theorem 1. *Let K be a $2m$ -dimensional Eulerian manifold. Then*

$$\chi(K) = \sum_{i=0}^{2m} \left(-\frac{1}{2}\right)^i f_i(K) \quad (1)$$

holds, where $f_i(K)$ is the number of i -simplices of K .

A finite simplicial complex L is called a *flag complex* if every collection of vertices of L which are pairwise adjacent spans a simplex of L . The formula (1) was proved in [1] under the additional assumptions that K is a PL-triangulation of a closed $2m$ -manifold and is a flag complex. M. W. Davis pointed out that the formula (1) follows from a result in [3], provided K is a flag complex (see *Note added in proof* in [1]). Both results follow from the considerations of the Euler characteristics of Coxeter groups. In this note, we deduce the formula (1) from the generalized Dehn-Sommerville equations proved by Klee [4].

Let K be a finite $(d-1)$ -dimensional simplicial complex and $f_i = f_i(K)$ the number of i -simplices of K as before. The d -tuple $(f_0, f_1, \dots, f_{d-1})$ is called the *f-vector* of K . The *f-polynomial* $f_K(t)$ of K is defined by

$$f_K(t) = t^d + f_0 t^{d-1} + \dots + f_{d-2} t + f_{d-1}.$$

Define the *h-polynomial* $h_K(t)$ of K ,

$$h_K(t) = h_0 t^d + h_1 t^{d-1} + \dots + h_{d-1} t + h_d,$$

Partially supported by the Grant-in-Aid for Scientific Research (C) (No.17560054) from the Japan Society for Promotion of Sciences.

by the rule $h_K(t) = f_K(t-1)$. The $(d+1)$ -tuple $(h_0, h_1, \dots, h_d)$ is called the h -vector of K . The h -vector of K satisfies the generalized Dehn-Sommerville equations, as stated below in Theorem 2.

Theorem 2 ([4]). *Let K be a $(d-1)$ -dimensional Eulerian manifold. Then*

$$h_{d-i} - h_i = (-1)^i \binom{d}{i} (\chi(K) - \chi(S^{d-1}))$$

holds for all i ($0 \leq i \leq d$).

Remark. Klee stated the generalized Dehn-Sommerville equations in terms of the f -vector rather than the h -vector. The formulae quoted in Theorem 2 are equivalent to those in [4] and can be found in [5]. Theorem 2 was also proved in [2] by a quite different method, provided that K is a triangulation of a closed manifold.

Now we prove Theorem 1. We have

$$h_K(-1) = \sum_{i=0}^{2m+1} (-1)^{2m+1-i} h_i = \sum_{i=0}^m (-1)^i (h_{2m+1-i} - h_i).$$

Now Theorem 2 asserts that

$$h_{2m+1-i} - h_i = (-1)^i \binom{2m+1}{i} (\chi(K) - 2).$$

Hence we obtain

$$h_K(-1) = (\chi(K) - 2) \sum_{i=0}^m \binom{2m+1}{i} = 2^{2m} (\chi(K) - 2). \quad (2)$$

On the other hand, we have

$$f_K(-2) = (-2)^{2m+1} + \sum_{i=0}^{2m} (-2)^{2m-i} f_i = 2^{2m} \left(-2 + \sum_{i=0}^{2m} \left(-\frac{1}{2} \right)^i f_i \right). \quad (3)$$

Since $h_K(-1) = f_K(-2)$ by the definition of the h -polynomial $h_K(t)$, Theorem 1 follows from (2) and (3).

REFERENCES

1. T. Akita, Euler characteristics of Coxeter groups, PL-triangulations of closed manifolds, and cohomology of subgroups of Artin groups, *J. London Math. Soc.* (2) 61 (2000), 721–736.
2. V. M. Buchstaber, T. E. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series 24, American Mathematical Society, Providence, 2002.
3. R. Charney, M. W. Davis, Reciprocity of growth functions of Coxeter groups, *Geom. Dedicata* 39 (1991), 373–378.
4. V. Klee, A combinatorial analogue of Poincaré’s duality theorem, *Canad. J. Math.* 16 (1964), 517–531.
5. E. Swartz, From spheres to manifolds, preprint (2005).

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO, 060-0810 JAPAN

E-mail address: akita@math.sci.hokudai.ac.jp